



A minimally invasive treatment
for stress urinary incontinence



This booklet provides information about the Bulkamid hydrogel for the treatment of stress urinary incontinence



BULKAMID[®]

What is stress urinary incontinence?

Stress urinary incontinence is the unintentional passing of urine during activity or exertion. It is caused by a weakness of the pelvic floor muscles

Stress urinary incontinence is common affecting an estimated 30% of women worldwide¹ and can have a significant impact on daily life affecting activities, relationships and emotional well-being². It can occur at any stage of life, with risk factors including pelvic disorders from childbirth, pelvic surgery and ageing




How is stress urinary incontinence normally treated?

Stress urinary incontinence is a very treatable condition. Doctors will advise of the different options available. These may include:

- o Pelvic muscle strengthening
- o Medications
- o Urethral bulking
- o Surgery e.g. mid-urethral tapes or colposuspension

What is urethral bulking?

Urethral bulking is the injection of a bulking agent, like Bulkamid, into the urethra to add volume to the tissue. This supports the closing mechanism of the urethra and provides better control of urine when you cough, laugh, exercise or change position



Urethral bulking offers a less invasive alternative to other SUI procedures (e.g. placement of mid-urethral slings), with fewer complications and a quicker recovery time³

What is Bulkamid?

Bulkamid is a soft gel consisting of 97.5% water and 2.5% polyacrylamide

It has been shown to maintain its shape and volume years after implantation, providing long term relief from urine leakage

How successful is Bulkamid?

Over three quarters of women who receive Bulkamid report to be cured or significantly improved³ and recent data has shown the effect can last up to 7 years⁴

If the effect of Bulkamid is insufficient following the first injection a top-up injection can be given to help achieve satisfactory resolution of symptoms

Is Bulkamid a safe procedure?

Bulkamid has treated over 80,000 women with stress urinary incontinence in 25 countries over 10 years. During that time few complications have been reported and there have been no long term complications. However, as with any invasive procedure complications may occur

Although rare, complications can include pain related to the procedure, a small amount of blood in the urine, delayed urination, painful urination, and / or urinary tract infection. These complications are all temporary and if they occur should resolve within a few days



What can I expect on the day of my operation?

Bulkamid is a minimally invasive procedure which will be completed in approximately 15-20 minutes. The procedure will usually take place in day surgery or outpatients' clinic and patients will normally be able to go home on the same day

Prior to the procedure patients will discuss whether to have a local or general anaesthetic with their doctor. Most patients who have undergone a Bulkamid procedure under local anaesthetic feel no more than a slight scratch as the needle enters the urethral wall

Once the procedure starts 3 or 4 deposits of Bulkamid (1.5 – 2ml total volume) will be placed into the urethral wall until coaptation of the urethral lumen is achieved

A patient should be able to return to normal activities within 24 hours subject to their doctor's advice

For further information please visit www.bulkamid.com

THE IDEAL BULKING SOLUTION

Laugh
Sneeze
Lift
Run
Cough
Jump



Speciality European Pharma Limited
14 Took's Court, London EC4A 1LB
Tel: +44 (0)20 7421 7400, Fax: +44 (0)20 7421 7401
www.specialityeuropeanpharma.com



Contura International A/S
Sydmarken 23, DK-2860 Soeborg
Tel: +45 81 100 900, Fax: +45 81 100 901
www.bulkamid.com, orders@bulkamid.com

January 2017

References

1. 5th International Consultation on Incontinence, Paris February 2012
2. Sinclair A, Ramsay I. Review: The psychological impact of urinary incontinence in women
3. Sokol ER, Karram MM, Dmochowski R. Efficacy and safety of polyacrylamide hydrogel for the treatment of female stress incontinence: a randomized, prospective, multicenter North American study. J Urol. 2014 Sep;192(3):843-9.
4. Lobodasch K & Brosche V. Long-term effectiveness and durability of Bulkamid® as primary treatment of stress urinary incontinence – a longitudinal study. ICS 2015 abstract.